Topic: Circumference and Area

Topic/Skill	Definition/Tips	Example
1. Circle	A circle is the locus of all points equidistant from a central point.	
2. Parts of a Circle	**Radius** – the **distance** from the **centre** of a circle to the **edge** **Diameter** – the total **distance** across the **width** of a circle **through the centre**. **Circumference** – the **total distance** around the **outside** of a circle **Chord** – a **straight line** whose **end points lie on a circle** **Tangent** – a **straight line** which **touches** a circle at exactly **one point** **Arc** – a **part of the circumference** of a circle **Sector** – the **region** of a circle enclosed by **two radii** and their intercepted **arc** **Segment** – the **region** bounded by a **chord** and the **arc** created by the chord	
3. Area of a Circle	$\boldsymbol{A = \pi r^2}$ which means 'pi x radius squared'.	If the radius was 5cm, then: $A = \pi \times 5^2 = 78.5cm^2$
4. Circumference of a Circle	$\boldsymbol{C = \pi d}$ which means 'pi x diameter'	If the radius was 5cm, then: $C = \pi \times 10 = 31.4cm$
5. π ('pi')	Pi is the circumference of a circle divided by the diameter. $\boldsymbol{\pi \approx 3.14}$	
6. Volume of a Cylinder	$\boldsymbol{V = \pi r^2 h}$	 $V = \pi(4)(5)$ $= 62.8cm^3$
7. Volume of a Prism	$\boldsymbol{V = Area\ of\ Cross\ Section \times Length}$ $\boldsymbol{V = A \times L}$	

8. Pythagoras' Theorem	For any **right angled triangle**: $$a^2 + b^2 = c^2$$  Used to find **missing lengths**. a and b are the shorter sides, c is the **hypotenuse** (**longest side**).	 $a = y,\ b = 8,\ c = 10$ $a^2 = c^2 - b^2$ $y^2 = 100 - 64$ $y^2 = 36$ $y = 6$
9. Direct Proportion	If two quantities are in direct proportion, **as one increases**, the **other increases** by the **same percentage**. If y is directly proportional to x, this can be written as $\boldsymbol{y \propto x}$ An equation of the form $\boldsymbol{y = kx}$ represents direct proportion, where k **is the constant of proportionality**.	
10. Real Life Graphs	Graphs that are supposed to model some real-life situation. The actual meaning of the values depends on the labels and units on each axis. The **gradient** might have a contextual meaning. The **y-intercept** might have a contextual meaning. The **area** under the graph might have a contextual meaning.	 A graph showing the cost of hiring a ladder for various numbers of days. The gradient shows the cost per day. It costs £3/day to hire the ladder. The y-intercept shows the additional cost/deposit/fixed charge (something

		not linked to how long the ladder is hired for). The additional cost is £7.
11. Conversion Graph	A line graph to **convert one unit to another**. Can be used to convert units (eg. miles and kilometres) or currencies ($ and £) Find the value you know on one axis, read up/across to the conversion line and read the equivalent value from the other axis.	 $8\ km = 5\ miles$
12. Depth of Water in Containers	Graphs can be used to show how the depth of water changes as different shaped containers are filled with water at a constant rate.	
13. Distance-Time Graphs	You can find the **speed** from the **gradient** of the line (Distance ÷ Time) The steeper the line, the quicker the speed. A **horizontal** line means the object is not moving (**stationary**).